

Digoxin (DIGO)

Source:

Cerba Lancet Africa

View our website:

cerbalancetafrica.com

Date & Time of Export:

15 April 2026 06:04



[Click here to view the results online](#)

TEST OVERVIEW

Test Name	Digoxin
Test Code	DIGO
Short Description	Digoxin

OVERVIEW

Test Name	Digoxin
Test Code	DIGO
Category	Biochemistry
TAT	Main Lab: 6 Hour(s) Family Site: <8hrs
Specimen(s)	1 x Venous blood - 5 mL Tube - Gold - SST-Serum Separator Tube

SPECIMEN(S)

SST-Serum Separator Tube

Specimen Type	SST-Serum Separator Tube
Specimen Format	Tube
Specimen Colour	Gold
Specimen Volume	5 mL
Sampling Order	2
Origin	Venous blood
Collection time after baseline	-
Transport Temperature	2-8°C
Accepted Other Specimens	Lithium Heparin Plasma Sodium Heparin Plasma EDTA Plasma

	Serum
TAT	Main Lab: 6 Hour(s) Family Site: <8hrs
Test Stability	Room Temp: 6 Hour(s) 2–8°C: 2 Day(s)

PATIENT INFORMATION

Clinical Information Required	Monitoring as part of treatment Drug name Frequency Date of last medication taken Time of last medication taken
Patient Collection Notes	Samples should be drawn 12 hours after oral administration of digoxin or 6 hours after IV administration It is recommended that a standardised 12-hour post-dose lithium concentration be used to assess adequate therapy. Peak concentration is reached 2 to 4 hours after an oral dose.

COMMENTS & NOTES

LOINC Code	535-3, 10535-3
Outwork	<input type="button" value="No"/>